

# University of Notre Dame Calculus III

## LECTURE 6: SPACE CURVES

---

### Vector Valued Functions and Space Curves

A vector-valued function is a function whose output is a vector. We have already encountered one: the vector equation for a line

$$\vec{l}(t) = \vec{P}_0 + t\vec{v}$$

More generally, they will have the form

$$\begin{aligned}\vec{r}(t) &= \langle f(t), g(t), h(t) \rangle \\ &= f(t)\hat{i} + g(t)\hat{j} + h(t)\hat{k}\end{aligned}$$

The input variable (in this case it's  $t$ ) is called the parameter. Since  $\vec{r}$  is a function, we can ask about its domain. The domain of a vector-valued function is the "intersection\*" of the domains of its components functions, that is, the values common to the domains of each of  $f$ ,  $g$ , and  $h$ .

**Example 1.** What is the domain of  $\vec{r}(t) = \langle \sqrt{4-t^2}, e^{-3t}, \ln(t+1) \rangle$ ?

**Solution:**

As with normal functions, we can take limits of vector-valued functions:

If  $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$ , then

$$\lim_{t \rightarrow a} \vec{r}(t) = \langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \rangle$$

And this leads us to the definition of continuity for vector valued functions:

**Definition 1.** A vector valued function  $\vec{r}(t)$  is called continuous at a if

$$\lim_{t \rightarrow a} \vec{r}(t) = \vec{r}(a)$$

Let's look at some examples of vector-valued functions.

**Example 2.**

- i)  $\vec{r}(t) = \langle \sin t, t \rangle$       ii)  $\vec{r}(t) = \langle \sqrt{2} \cos t, \sin t \rangle$   
iii)  $\vec{r}(t) = \langle \cos t, \sin t, t \rangle$       iv)  $\langle t, \sin t, 2 \cos t \rangle$   
v)  $\vec{r}(t) = \langle t \cos t, t \sin t, t \rangle$

**Solution:**

Often of importance is the intersection\* of surfaces. The result is generically a curve (i.e., the intersection\* of two planes is a line, as we know). It is important to be able to quantify these "curves of intersection\*" by parameterizing them (finding a vector-valued function whose image is the curve).

**Example 3.** Find a vector function representing the intersection\* of  $z = x^2$  and  $x^2 + y^2 = 4$ .

**Solution:**

### Extra Examples

1. Sketch the curve given by  $\mathbf{r}(t) = \langle \sin 2\pi t, t, \cos 2\pi t \rangle$ , indicating by an arrow the direction in which  $t$  increases.
2. What is the domain of the vector function  $\mathbf{r}(t) = \langle \frac{1}{t+2}, \cos t, \ln(16 - t^2) \rangle$ ?
3. Determine the parametric equations and the vector equation for the intersection\* of the plane  $y + z = 2$  and the cylinder  $x^2 + y^2 = 1$ .