

# University of Notre Dame Calculus III

## LECTURE 26: THE FUNDAMENTAL THEOREM OF LINE INTEGRALS

---

Throughout the rest of the class we will be generalizing the result so fundamental, it says so in its name: the fundamental theorem of calculus

$$\int_a^b \frac{d}{dx} f(x) dx = f(b) - f(a)$$

There will be a theme to each of these we will be trading derivatives for boundaries (or vice versa). Notice  $\partial[a, b] = \{a, b\}$ .

First we need some terminology.

### Conservative Vector Fields and Potentials

If  $\vec{F} = \langle P, Q \rangle$  is conservative, then  $P = f_x$  and  $Q = f_y$  for some  $f = f(x, y)$ . So, by Clairaut's theorem  $P_y = Q_x$  (as long as  $\vec{F}$  is  $C^1$ ). This gives us a test for conservative vector fields. Let  $\vec{F}$  be a  $C^1$  vector field.

(a) If  $\vec{F}(x, y) = \langle P, Q \rangle$  and  $\vec{F}$  is conservative, then

$$P_y = Q_x$$

(b) If  $\vec{F}(x, y, z) = \langle P, Q, R \rangle$  and  $\vec{F}$  is conservative, then

$$\text{curl } \vec{F} = \vec{0}$$

**Note:** We don't know what curl is yet, but will learn shortly.

These are tests we can use to determine whether  $\vec{F}$  is not conservative. If  $P_y \neq Q_x$  in (a) or  $\text{curl } \vec{F} \neq \vec{0}$  in (b), then  $\vec{F}$  is not conservative. If the equations hold then we don't know if  $\vec{F}$  is conservative or not. A curiosity would be to know when these results are reversible, i.e., when  $P_y = Q_x$  or  $\text{curl } \vec{F} = \vec{0}$  implies  $\vec{F}$  is conservative. To answer this, we need two definitions.

**Definition 1.** A curve which does not cross itself anywhere between its endpoints is called simple.

**Definition 2.** A region  $D$  is called simply connected if  $D$  is connected and every simple closed curve in  $D$  encloses only points in  $D$ .

Here are the reversals:

**Theorem 3.** a) If  $\vec{F} = \langle P, Q \rangle$  is a  $C^1$  vector field on an open, simply connected region  $D$ , and  $P_y = Q_x$ , we have that  $\vec{F}$  is conservative.

b) If  $\vec{F} = \langle P, Q, R \rangle$  is a  $C^1$  vector field on all of  $\mathbb{R}^3$  and  $\text{curl } \vec{F} = \vec{0}$ , then  $\vec{F}$  is conservative.

Often, the easiest way to determine whether a vector field is conservative is to just try to find a potential.

## The Fundamental Theorem of Line Integrals

**Theorem 4.** *Fundamental Theorem of Line Integrals* Let  $C$  be a smooth curve given by  $\vec{r}(t)$ ,  $a \leq t \leq b$ . Let  $f$  be a differentiable function of 2 or 3 variables whose gradient is continuous on  $C$ . Then

$$\int_C \nabla f \cdot d\vec{r} = f(\vec{r}(b)) - f(\vec{r}(a))$$

*Proof.*

$$\nabla f = \langle f_x, f_y, f_z \rangle$$

$$\begin{aligned} \int_C \nabla f \cdot d\vec{r} &= \int_a^b \nabla f(\vec{r}(t)) \cdot \vec{r}'(t) dt = \int_a^b \frac{d}{dt} f(\vec{r}(t)) dt \\ &= f(\vec{r}(b)) - f(\vec{r}(a)) \end{aligned}$$

□

**Definition 5.** We say a line integral  $\int_C \vec{F} \cdot d\vec{r}$  is independent of path if given any two curves  $C_1$  and  $C_2$  in the domain of  $\vec{F}$  which start and end at the same place we have

$$\int_{C_1} \vec{F} \cdot d\vec{r} = \int_{C_2} \vec{F} \cdot d\vec{r}$$

The theorem tells us then that line integrals of conservative vector fields are independent of path.

**Definition 6.** A curve is closed if it starts and ends at the same point.

**Theorem 7.**  $\int_C \vec{F} \cdot d\vec{r}$  is independent of path in  $D$  if and only if  $\int_C \vec{F} \cdot d\vec{r} = 0$  for all closed paths  $C$  in  $D$ .

**Definition 8.** A set  $D$  is open if for every point  $P$  in  $D$ , we can fit a disk of radius  $\epsilon > 0$  ( $\epsilon$  as small as needed) around  $P$  inside  $D$ .

**Definition 9.** A set  $D$  is connected if any two points in  $D$  can be joined by a path in  $D$ .

**Theorem 10.** Suppose  $\vec{F}$  is a vector field which is continuous on an open and connected set  $D$ . If  $\int_C \vec{F} \cdot d\vec{r}$  is independent of path in  $D$ , then  $\vec{F}$  is conservative on  $D$ .

**Example 1.** Compute  $\int_C (\ln y + 2xy^3) dx + (3x^2y^2 + \frac{x}{y}) dy$  where  $C$  has parametric equations

$$x = \frac{1}{2}t^2 + 2, \quad y = e^t(1 + 2t - t^2), \quad 0 \leq t \leq 2.$$

**Solution:**

Let's do an application to end this section: Let  $\vec{F}$  be a continuous force field which moves an object along a path  $C$  given by  $\vec{r}(t)$ ,  $a \leq t \leq b$  where  $\vec{r}(a) = A$  and  $\vec{r}(b) = B$ . By Newton's second law

$$\vec{F}(\vec{r}(t)) = m\vec{r}''(t)$$

Thus, the work done by  $\vec{F}$  is then

$$\begin{aligned} W &= \int_C \vec{F} \cdot d\vec{r} = \int_a^b \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt = \int_a^b m\vec{r}''(t) \cdot \vec{r}'(t) dt \\ &= \frac{m}{2} \int_a^b \frac{d}{dt} [\vec{r}'(t) \cdot \vec{r}'(t)] dt = \frac{m}{2} \int_a^b \frac{d}{dt} \|\vec{r}'(t)\|^2 dt \\ &= \frac{1}{2} m \|\vec{r}'(t)\|^2 \Big|_a^b = \frac{1}{2} m \|\vec{r}'(b)\|^2 - \frac{1}{2} m \|\vec{r}'(a)\|^2 \\ &= \frac{1}{2} m (\|\vec{v}(b)\|^2 - \|\vec{v}(a)\|^2) \end{aligned}$$

The quantity  $\frac{1}{2} m (\|\vec{v}(t)\|)^2$  is the kinetic energy at time  $t$ , so we can write  $W = K(B) - K(A)$ . Now, let's assume that the force  $\vec{F}$  is conservative. Then  $\vec{F} = \nabla f$ . Typically, in Physics, the potential energy is defined as  $P = -f$  so that  $\vec{F} = -\nabla P$ . Then, the FToLI gives

$$W = \int_C \vec{F} \cdot d\vec{r} = - \int_A^B \nabla P \cdot d\vec{r} = -(P(B) - P(A)) = P(A) - P(B)$$

Combining this with what we know about kinetic energy we have

$$K(B) - K(A) = W = P(A) - P(B) \quad \iff \quad P(A) + K(A) = P(B) + K(B)$$

This is the Law of Conservation of Energy.

Let's do an example of finding a potential for a 3D vector field.

**Example 2.** Determine whether the vector field

$$\vec{F} = \langle e^x \sin yz, ze^x \cos yz, ye^x \cos yz + 3z^2 \rangle$$

is conservative. If so, find a potential.

Then find

$$\int_C \vec{F} \cdot d\vec{r}$$

where  $C$  is a smooth curve where given by  $\vec{r}(t)$  that starts at  $\vec{r}(0) = \langle 0, \pi/2, 1 \rangle$  and ends at  $\vec{r}(\pi) = \langle 1, 0, 2 \rangle$ .

**Solution:**